
SUPPLEMENTAL LABEL



Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, NC 27419
www.syngenta-us.com

QUILT®

Fungicide

EPA Reg. No. 100-1178

Supplemental label for aerial application on corn

Active Ingredient:	
Azoxystrobin	7.0%
Propiconazole	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Quilt as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling supersedes the Directions for Use on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

APPLICATION PROCEDURES

Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner that will result in exposure to humans or animals.

Ground Application:

- For field crops (non-trees), apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 50 gallons of water per acre unless specified otherwise.

Aerial Application:

- Use only on crops where aerial applications are indicated.
- For field crops (non-trees), apply in a minimum of 2 gallons of water per acre unless specified otherwise.
- For tree crops, apply in a minimum of 10 gallons of water per acre unless specified otherwise.
- Do not apply via ULV.

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Corn Field Pop (Includes Seed Production)	Northern corn leaf blight (<i>Setosphaeria turcica</i>) Northern corn leaf spot (<i>Cochliobolus carbonum</i>) Southern corn leaf blight (<i>Cochliobolus heterostrophus</i>) also known as Helminthosporium leaf blights (<i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i>)	7 - 14	For leaf blights apply Quilt when disease first appears. Continue on a 7 to 14 day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate. Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.
	Rusts (<i>Puccinia</i> spp.) Gray leafspot (<i>Cercospora zeae-maydis</i>) Eye spot (<i>Aureobasidium zeae</i>)	10.5 - 14	Apply Quilt when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule. Apply no more than 2 applications of Quilt or any other Group 11 fungicide per year.
Application Instructions: Quilt is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume should be used to provide thorough coverage. Quilt may be applied by ground, air, or chemigation.			
Specific Use Restrictions: <ol style="list-style-type: none"> 1) Do not apply within 30 days of harvest for forage, grain, or stover. 2) Do not apply more than 56 fl. oz./A/season of Quilt. 3) Do not apply more than 28 fl. oz. (0.224 lb. a.i. propiconazole) for field corn harvested for forage. 4) Do not apply more than 0.45 lb. a.i. propiconazole containing products/A/season. 5) Do not apply more than 2.0 lbs. a.i. azoxystrobin containing products/A/season. 			

Quilt® and the Syngenta logo are trademarks of a Syngenta Group Company

© 2007 Syngenta
 Syngenta Crop Protection, Inc.
 Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1178A-S7 0507

QUI 1178A-S7 0507 – df – 5/25/07